

**Review: *Vaccination Roulette: Experiences, risks, and alternatives*,
Australian Vaccination Network, Bangalow, NSW, 1998**

Pages: 81

Shocking Revelations

This is a wide-ranging look at the lack of vaccine safety.

It details the disastrous history from smallpox and polio vaccine creation (e.g. the Simian Virus 40 permanent genetic contamination of 400,000 vaccinees in 1954).

Also included is basic pathological information on the big ‘vaccine-preventable diseases’ (tetanus, pertussis, rubella, varicella, mumps, and measles).

Vaccine injury and common symptoms are listed in the context of the Australian medical profession’s woeful ignorance of such injuries (only 3% of doctors are even aware of the existence of ADRAC or AEFVSS [now replaced by ACSOM and ATAGI]).

While the material is shocking and negative to a degree, there are positive alternative health options like homeopathy and naturopathy.

Another piece of the puzzle is media malfeasance and propaganda in maligning the vaccine-injured and health advocates as ‘anti-vaxxers’.

Chapters are written by different contributors which each bring their own perspective and expertise.

Overall, a great attempt at making people medically informed about a risky and dangerous medical procedure marketed as Pharma’s panacea for the world’s problems. Though published in 1998, the refusal to do a true double-blind randomised placebo-controlled study of vaccinated versus unvaccinated, or even study vaccinated versus unvaccinated using the CDC’s comprehensive Vaccine Safety Datalink database means the challenge remains unanswered twenty years later.

Foreword (pp. viii-x)

Since exams must be passed, medical students believe all they are taught, and after graduation they are mainly educated by drug company representatives. Any adverse events to vaccination are assumed coincidental.

Introduction (pp. 1-8)

If vaccination gave its promised protection then no vaccinated person is exposed to the risk of the unvaccinated. By targeting the unvaccinated in order to protect the vaccinated, pro-vaccinators expose the weak underbelly of vaccines.

Compulsion and coercion like “money for jabs” is a denial of the right of choice.

The Australian Constitution, Section 51, Part 23A ‘civil conscription’ includes any compulsory medical procedure.

Forced vaccinators appeal to the “common good” as a rationale.

Public awareness is nil so as not to endanger pharma profits and the whole shaky foundation of vaccination policy.

I) Vaccine Alert: The Case for Caution Before Vaccinating (pp. 9-38)

Vaccination always involves risk and a two-month old baby is far more vulnerable. A baby’s brain is not fully developed till age five.

Vaccine-ingredient damage cannot be reversed.

Vaccines don’t give long term protection, they severely suppress the immune system (which is why sick children should not be vaccinated), and they can never eradicate a disease (e.g. smallpox has enormous non-human reservoirs).

Human DNA in vaccines may cause the body to develop anti-DNA antibodies causing lupus for example.

Multi live-virus vaccines may increase genetic recombination risk and new diseases.

Natural immunity is the only way for mothers to pass on antibodies to newborns via breast milk since babies do not yet have a functioning immune system. Mother-titer levels are about 1,000, or a fifth of what they were three generations ago.

The OPV can spread polio via the child's stool for up to eight weeks after vaccination.

III) Demyelination and Encephalitis (pp. 39-72)

The pertussis vaccine has been known to damage brain and nerves since 1920 and is itself an extremely strong adjuvant.

Allergic hypersensitisation makes the immune system sensitive to any and every allergen.

Injection of *Bordetella pertussis* into lab animals caused experimental allergic encephalopathy (EAE).

Encephalopathy of the nerve myelin makes it faulty or without sufficient insulation.

Vaccine damage may manifest as sleep disturbances, 'minimal' brain damage, dyslexia, hyperactivity, lethargy, bulimia, anorexia, celiac disease-nerve palsies, tics, muscle spasms, and developmental delay.

Myelin is a tough, white fatty waterproof substance covering neurons and axons after they are grown. Some nerves do not myelinate until eight months of age, and some continue up to age forty-five!

Brain demyelination degrades intellect, causes a loss of sense of humour, empathy, reading and calculational abilities.

Vaccines are not standardised between manufacturers or between batches. There is also no lot or batch numbers. Unless they are refrigerated, they lose potency and shelf life.

Pertussis toxin can alter the catecholaminergic and GABAergic systems of neurotransmitters. Normally it cannot cross the BBB, however, it can in the presence of high blood pressure, acute stress, seizures, and hypoxia.

Pertussis vaccine immunity begins waning after five years.

Whooping cough begins with a cold then develops into a cough, where the bacteria irritate and inflame the airways causing thick mucus to develop. There is a ten day to two-week window for treatment before coughing becomes serious.

Vitamin C treatment held infections to 20 days and prevented the spasmodic stage while the vaccinated suffered for 34 days.

After infection on milk or starch foods should be given at all.

Vaccinators keep 1:1000 epinephrine available in case of anaphylaxis.

Many doctors would go under without money from vaccine companies.

Sanitation improvement between 1850 and 1940 reduced disease incidence by 90% in Britain.

Live-attenuated viruses such as MMR may cause the disease, which is why immuno-compromised children must be kept away from the recently-vaccinated.

97.5% of measles cases are misdiagnosed.

The Hepatitis B vaccine was first marketed at drug users, prostitutes, prisoners, and homosexuals. Its effective duration is unknown. The virus is spread by blood or semen and is most prevalent in Asia and Africa (10% of the population), but in the West is 0.1% to 0.5%.

95% of hepatitis B infections will clear after four weeks of nausea and will give permanent immunity. The fatality rate is between 0.1% and 1.4%.

IV) “There Are Three Kinds of Lies: Lies, Damned Lies, and Statistics” **(pp. 73-82)**

The AVN addresses compulsory vaccination, indiscriminate mass vaccination, and the official refusal to acknowledge vaccine injuries.

The medical industrial complex succeeded by first gaining a monopoly over medical practice.

The MMR vaccine has over 600 side effects.

V) Proof of Vaccination Damage (pp. 83-131)

Dr Robert Mendelsohn: “Doctors in general should be treated with about the same degree of trust as used-car salesmen.”

In 1997, the AVN contacted almost 400 families of children either injured or killed by vaccines.

Only about 3% of doctors in Australia know about the Adverse Drug Reaction Advisory Committee (ADRAC) or the Adverse Events Following Vaccination Surveillance Scheme (AEFVSS).

Sweden stopped vaccinating against whooping cough in 1979.

The whole-cell pertussis vaccine is only 36-48% effective.

Dr Gavin Frost: “Doctors are busy people and reporting vaccine reactions is not their first priority.”

Over 79% of children were fully vaccinated during a 1994 Western Sydney measles outbreak.

There are at least 60,000 chemical compounds which share a basic benzene ring structure. Many are mutagenic or carcinogenic.

Allergies diminish the immune response and increase infection vulnerability.

Chronic arthritis is associated with natural rubella and is thought to be related to viral antigen isolated from body tissue.

Guillain-Barre and CFS are similar in symptoms.

Carbolic acid (phenol) is a sweet-smelling acid that affects the CNS, liver and kidney, and is readily absorbed through the skin. It can cause seizures, there is no antidote and one gram can be fatal.

20% of people are hypersensitive to formaldehyde.

Hg and allergens combine to rupture WBCs, it suppresses the immune system, causes scleroderma, heart disease, and destroys RBCs and damages chromosomes.

$\text{Al}(\text{OH})_3$ and AlPO_3 are common adjuvants.

Al changes neuron structure.

Chemical exposure may be acute, subacute, subchronic, or chronic and routes may be intraperitoneal, intravenous, subcutaneous, oral, or dermal absorption.

After sensitisation, allergic reactions can be triggered at any time.

Most nerve myelin is laid down before age five.

Jonas Salk and Albert Sabin worked concurrently on the polio vaccine. Since liquid isolation in 1908, no one knew how to keep the virus alive ex-vivo.

In 1948, Dr John Enders and two co-workers grew polio in cells from cast-away human placentas.

In 1955, Wendell Stanley crystallised the virus in his Berkeley lab.

In a 1954 field test, 400,000 were vaccinated causing over 200 polio cases and 11 deaths.

In another incident up to thirty Salk vaccines and Sabin sugar cubes were contaminated with Simian Virus 40 from Asian rhesus monkeys. Bernice Eddy discovered live monkey viruses in IPV in 1954 which the Merck vaccine division later confirmed.

Children of Salk-vaccinated children are at a thirteen-fold greater risk of brain tumours and German scientists have found SV40 in about 30% of all cases.

On May 23, 1935, Dr Harris, Minister of Public Health in Victoria said, "I would never allow a child of mine to be immunised."

Childhood-diphtheria diagnosis would often be revised as tonsillitis. Anti-diphtherium serum is made by injecting a horse with increasing 'toxin' doses then after several months blood is drawn from its neck, and the serum drained off.

The US government introduced forced diphtheria vaccination in 1945 into Germany.

Children with encephalitis are emotionally unstable, have reduced volitional inhibitions, and exhibit evening restlessness.

In 1988 in the US, 13% of children had 'minimal brain damage'.

VI) How Clean Are Vaccines (pp. 132-145)

Most vaccines use monkey kidneys with 20-80% discarded in the process due to contaminate concerns.

Other-virus contamination is unavoidable for live-virus vaccines.

Formalin does not kill SV40.

Measles vaccine now uses canine kidneys.

AIDS began in the Belgian Congo as a result of an SV40-WHO OPV vaccination campaign.

Companies spend \$8,000-\$10,000 per doctor p.a. on product marketing.

Japan introduced acellular pertussis vaccine in 1981.

The "cerebral cry" is due to brain inflammation.

Standard vaccination procedure should be:

1. Detail risk.
2. Prepare for any adverse reactions.
3. Read the product information.

4. Ask for and record consent.
5. Give a pre-vaccination questionnaire.
6. Check for contraindications.
7. Confirm patient identity.
8. Give paracetamol.
9. Check the vaccine product including prior storage conditions and expiry date.

VII) Vaccination: The Latest Sacrament of Medical Science (pp. 146-154)

The Government introduced the Australian Childhood Immunisation Register (ACIR) in 1996.

Doctors get increased patient consultation, provide treatment for vaccine-related illnesses, and counselling referrals for vaccine-refusing parents.

VIII) Are There Alternatives? (pp. 155-174)

Reflexology posits that the hands, ears, and feet replicate the whole body's condition.

A swollen fontanelle is where the flat bones of a baby's skull do not meet so any encephalitis causes membrane to bulge outside of it.

Samuel Hahnemann (1755-1843) founded Homeopathy. Its Bible is the Organon and it describes an energy or "vital force".

A *similum* is always a single remedy. and most remedies have come via trial and error or *provings*. Groups of healthy people take remedies and keep meticulous diaries, with all objective and subjective symptoms written up in a *Materia Medica*. Homeopaths assign disease as a result of Vital Force compromise, e.g. by overexposure to cold, wet or hot conditions, or emotional shock.

Tinctures are made mostly from plants and minerals.

Remedies are either X (decimal) or C (centesimal) scales. An X remedy combines one part tincture with nine parts ethanol (a C is one to ninety-nine). Progressive dilutions occur via shaking (“succussing”) the bottle.

Eczema moves down the trunk and torso towards the extremities before disappearing.

Children secrete polio virus post-vaccination up to six weeks.

The first medical book was written 300BC by Huang di Nei jing, *The Yellow Emperor’s Classic of Internal Medicine*.

The Chinese would blow powdered small pox scabs up the nose as an inoculation.

IX) Naturopathic Care of Childhood Infectious Diseases (pp. 175-194)

The Naturopath aims to bring the diseased part of the body into harmony with the whole.

Harold Sherman: “The more you depend on forces outside yourself, the more you are dominated by them.”

Primum non nocere.

Chickenpox (varicella) incubates 7-21 days and is infectious a few days prior to the rash until the last spot or blister has scabbed. It spreads via exhaled vapour droplets.

Rubella (Latin “reddish”) is spread by close contact and vapour droplets. It incubates 14-21 days and is infectious a week before and five days after the rash. This is the only known virus to damage babies in utero.

Measles incubates 8-21 days and is infectious 4 days before and 5-10 days after. A rash first appears behind the ears and frontal hairline.

Mumps incubates 12-28 days and is infectious 2 days before and 10 days after. Swelling starts in the salivary glands.

Whooping cough is caused by the *Bordetella pertussis* bacteria and incubates 7-21 days and is infectious up to 5 weeks after the first bout of

coughing starts. The Chinese call it the “100 day cough”. Protection isn’t transferred from mother to baby.

Tetanus is caused by the anaerobic *Clostridium tetani* bacteria which lives freely in soil, building dust and gardens. Oxygen is fatal to it. It incubates 6-10 days but possibly a month and causes body muscle tightening and jaw spasms.

X) Homeoprophylaxis (pp. 195-206)

Hib is Haemophilus influenza type b meningitis.

General protection comes from dietary, herbal, nutritional, chiropractic, homeopathic, and constitutional sources.

Disease-specific protection is from remedies using the Law of Similars to target diseases.

Doctors prescribe Ipecac to induce vomiting in stomach poisoning cases.

In 1801, Dr Hahnemann wrote “The Cure and Prevention of Scarlet Fever” in which he prescribed Belladonna as a treatment.

HP remedies are non-toxic and are 89% effective.

Germes can only exist on a medium of decaying organic matter, else they have nothing to feed on.

Toxaemia causes disease from malnourishment, poor diet, unhealthy living, overwork, etc.

XI) The Role of the Media in Vaccination Campaigns (pp. 207-220)

XII) Vaccine Safety and Effectiveness (pp. 221-257)

Of 987 doctors, only 30% used temperature monitors in their vaccine refrigerators, 26% knew the correct expiry time of the Sabin vaccine after opening (eight hours), and 41% correctly injected infants in the anterolateral aspect of the thigh.

Against a DT control (!), DTP was found to be 84% efficacious.

Of 53 SIDS deaths between 1979 and 1980 in LA, 27 had received a DTP vaccine within the previous 28 days.

SV40 was identified in DTP vaccines by reverse transcriptase analyses up to 1985.

Measles virus may persist in intestinal tissue, particularly Chron's disease sufferers.

Radiculitis is spinal column nerve root inflammation.

Strabismus is eye-squinting.

Varivax vaccinees may still transmit chickenpox virus. Women should not get pregnant within three months of vaccination, and no one would use salicylates because it may cause Reyes syndrome (liver and brain swelling). This vaccine was cultured in lung tissue from two human aborted babies.

In Sydney in the 1980s, only 50% of 7-12 year old full-vaccinated children still had antibodies to all three polio strains.

Vaccine damage in children leads to marriage breakdown (the father usually leaves), drug and alcohol dependency and distress (as there is no time for other family members), as well as mother guilt.

In 1984, T-helper lymphocytes dropped in 11 routine tetanus vaccinees. This is also associated with AIDS.

The Salk IPV does not harbour SV40.

Pertussis causes about 600,000 deaths p.a. worldwide.

XIII) Enlisting the Law to Fight Discrimination (pp. 258-293)

XIV) To Vaccinate or Not? The Choice Is Yours (pp. 295-307)

The Rubella vaccine is called RA27/3 since the virus was isolated from the 27th aborted baby at the Wistar Institute in the 1964 outbreak.

XV) Is Vaccination a Game of Roulette (pp. 309-326)

Roulette is A French game from the 17thC played on a *roulette* (a spinning horizontal wheel).